

1. PaDIL Species Factsheet



Scientific Name:

Xiphinema americanum (Cobb, 1913) Micoletzky, 1922
(Nematoda: Dorylaimida: Longidoridae)

Common Name

Dagger Nematode

Live link: <http://www.padil.gov.au/maf-border/Pest/Main/141265>

Image Library

New Zealand Biosecurity

Live link: <http://www.padil.gov.au/maf-border/>

Partners for New Zealand Biosecurity image library



Manaaki Whenua
Landcare Research

Landcare Research — Manaaki Whenua

<http://www.landcareresearch.co.nz/>

Biosecurity New Zealand

Tiakitanga Pūtaiao Aotearoa

MPI (Ministry for Primary Industries)

<http://www.biosecurity.govt.nz/>

2. Species Information

2.1. Details

Specimen Contact: MAF Plant Health & Environment Laboratory (Nematode collection held by Zeng Zhao at Landcare Research) - PHEL_Entomology@maf.govt.nz

Author: Zhao, Z. Q. & Crosby, T. K.

Citation: Zhao, Z. Q. & Crosby, T. K. (2011) Dagger Nematode (*Xiphinema americanum*) Updated on 7/7/2011
Available online: PaDIL - <http://www.padil.gov.au>

Image Use: Free for use under the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0)

2.2. URL

Live link: <http://www.padil.gov.au/maf-border/Pest/Main/141265>

2.3. Facets

Commodity Overview: Field Crops and Pastures, Forestry, Horticulture

Commodity Type: 1 Other, Basil and Mint, Citrus produce, Coconut, Coffee beans & Kape, Fabaceous produce, Figs & Breadfruit, Mango, Nuts, Rosaceous produce, Tomato, Tamarillo & Egg plant

Groups: Nematodes / Roundworms

Status: NZ - Exotic

Pest Status: 0 Unknown

Distribution: Afrotropic, Australasia, Indo-Malaya, Nearctic, Neotropic, Palearctic

Host Family: Anacardiaceae, Apiaceae, Apocynaceae, Arecaceae, Asteraceae, Caryophyllaceae, Cornaceae, Cucurbitaceae, Ericaceae, Euphorbiaceae, Fabaceae, Fagaceae, Lamiaceae, Malvaceae, Moraceae, Myrtaceae, Oleaceae, Pinaceae, Pittosporaceae, Poaceae, Rosaceae, Rubiaceae, Rutaceae, Salicaceae, Sapindaceae, Sapotaceae, Solanaceae, Ulmaceae, Verbenaceae, Vitaceae, Aceraceae, Chenopodiaceae, Dipterocarpaceae, Linaceae, Plantaginaceae, Tiliaceae

2.4. Other Names

American Dagger Nematode

Tobacco Ring Spot Nematode

Tylencholaimus americanus (Cobb, 1913) Micoletzky, 1922

Xiphinema californicum Lamberti & Bleve-Zaches, 1979

Xiphinema taylori Lamberti et al., 1991

2.5. Diagnostic Notes

Diagnostic characters: (Dr Zeng Qi Zhao, Landcare Research, Auckland, New Zealand).


These nematodes are minute (the largest species of the group being 2.2 mm in length), soft-bodied, vermiform, nearly transparent animals. They have a hard, needle-like 'stylet' (odontostyle and odontophore) at the mouth-end of the body which is capable of being extruded to puncture plant cells. The member species of the *X. americanum* group can be readily distinguished from other *Xiphinema* spp. by the following characteristics: small body length, relatively short (usually <150 µm) stylet (odontostyle + odontophore), thick cuticular lining of the *Xiphinema americanum* sensu lato pharynx, males usually absent, female genital branches equally developed, uterus short and without Z-organ, presence of symbiotic bacteria in the oocytes

and in the intestines of juveniles, short conoid tail with rounded terminus, females without sperm, males with posteriormost medioventral supplement close to the paired precloacal papillae.

The separation of the individual species within the group is difficult and still subject to controversy among specialists. Lamberti & Bleve-Zacheo (1979) provide details of many of the species, and Ebsary et al. (1989) provide a key to the species occurring in North America.

Plants whose roots are being attacked by *X. americanum* sensu lato, in the absence of virus, generally show no clear characteristic symptoms in the aerial parts. With high populations, a general reduction in vigour is observed and this appears in characteristic patches in the crop corresponding to the highest concentration of nematodes. Under heavy attack, the roots show swellings close to the root tips. When nematode feeding results in virus transmission, the characteristic symptoms of the particular virus in the crop concerned develop. These usually first appear in the aerial parts of the plant in the growing season after transmission to the roots has occurred.


3. Diagnostic Images




Slide mounted
Anterior - Female: Zeng Zhao Landcare Research



Slide mounted
Full Body Image - Female: Zeng Zhao Landcare Research



Slide mounted
Pharyngeal Region - Female: Zeng Zhao Landcare Research



Slide mounted
Pharyngeal Region - Female: Zeng Zhao Landcare Research



Slide mounted
Pharyngeal Region - Female: Zeng Zhao Landcare Research



Slide mounted
Reproductive Region Vulva - Female: Zeng Zhao Landcare Research

Results Generated:

Sunday, October 25, 2020

